

REVIEW

Xanthine oxidoreductase in cancer: more than a differentiation marker

Maria Giulia Battelli, Letizia Polito, Massimo Bortolotti & Andrea Bolognesi

Department of Experimental, Diagnostic and Specialty Medicine – DIMES, Alma Mater Studiorum – University of Bologna, General Pathology Unit, Via S. Giacomo 14, 40126 Bologna, Italy

Keywords

Differentiation, oncogenesis, reactive oxygen and nitrogen species, uric acid, xanthine oxidoreductase

CorrespondenceLetizia Polito, Department of Experimental, Diagnostic and Specialty Medicine – DIMES, Alma Mater Studiorum – University of Bologna, General Pathology Unit, Via S. Giacomo 14, 40126 Bologna, Italy.
Tel: +39 051 2094700; Fax: +39 051 2094746; E-mail: letizia.polito@unibo.it**Funding Information**

This work was supported by the Pallotti Legacies for Cancer Research. No relationship is present regarding any organization or entity having a direct financial or personal interest in the subject matter or materials discussed in the article.

Received: 4 September 2015; Revised: 5 November 2015; Accepted: 8 November 2015

Cancer Medicine 2016; 5(3):546–557

doi: 10.1002/cam4.601

Introduction: Biology of Xanthine Oxidoreductase

Xanthine oxidoreductase catalyzes the oxidation of hypoxanthine to xanthine and of xanthine to uric acid, that is, the last two steps of purine catabolism in the highest uricotelic primates. In addition, XOR can oxidize different endogenous metabolites as well as various xenobiotics, including toxic substances and anticancer drugs (reviewed in [1]). By generating irreversible products, XOR has a rate-limiting role in purine metabolism because its action precludes the salvage pathway of purine nucleotides (Fig. 1).

Abstract

Human xanthine oxidoreductase (XOR) catalyzes the last two steps of purine catabolism and is present in two interconvertible forms, which may utilize O₂ or NAD⁺ as electron acceptors. In addition to uric acid, XOR products may comprise reactive oxygen and nitrogen species that have many biologic effects, including inflammation, endothelial dysfunction, and cytotoxicity, as well as mutagenesis and induction of proliferation. XOR is strictly modulated at the transcriptional and post-translational levels, and its expression and activity are highly variable in cancer. Xanthine oxidoreductase (XOR) expression has been negatively associated with a high malignancy grade and a worse prognosis in neoplasms of the breast, liver, gastrointestinal tract, and kidney, which normally express a high level of XOR protein. However, the level of XOR expression may be associated with a worse outcome in cancer of low XOR-expressing cells, in relation to the inflammatory response elicited through the tissue damage induced by tumor growth. Xanthine oxidoreductase (XOR) has been implicated in the process of oncogenesis either directly because it is able to catalyze the metabolic activation of carcinogenic substances or indirectly through the action of XOR-derived reactive oxygen and nitrogen species. The role of uric acid is characterized by both oxidant and antioxidant action; thus, it is still debatable whether control of uricemia may be helpful to improve the outcomes of tumor illness.

Xanthine oxidoreductase is a homodimeric metalloflavoprotein with a molecular mass of approximately 300 kDa. Each subunit includes a molybdenum-containing molybdopterin cofactor, two nonidentical iron–sulfur redox centers, and one flavin adenine dinucleotide (FAD) cofactor (reviewed in [2]). Low levels of XOR activity are almost ubiquitous in mammalian tissue, with the highest levels present in the liver, intestine, kidney, lactating mammary gland, and vascular endothelial cells (reviewed in [3]).

Xanthine oxidoreductase is strictly regulated at the transcriptional and post-translational levels. The expression of XOR protein is increased by various hormones, growth factors, inflammatory cytokines, irritative stimuli, and low




Figure 1. Substrates of Xanthine oxidoreductase (XOR) activity. (A) XOR catalyzes the last two steps of purine catabolism and generates xanthine and uric acid, therefore precluding the salvage pathway of purine nucleotides. (B) XOR can oxidize different endogenous metabolites, including NADH and nitrate, thus producing reactive species of both oxygen and nitrogen. XOR can also oxidize various xenobiotics, hence activating both carcinogens and anticancer drugs.

oxygen tension. The post-translational modulation of XOR consists of quantitative and qualitative changes in its activity. Interconvertible demolybdo- and/or desulfo-forms of XOR protein are known to be inactive in xanthine catalysis, but they may still oxidize NADH. XOR activity may also be controlled through the switch between XOR dehydrogenase and oxidase forms. Such conversion may occur in different pathologic circumstances, including tissue damage by physical agents, poisoning with endogenous or toxic substances, and various hypoxic/ischemic conditions (reviewed in [4]).

Xanthine oxidoreductase is constitutively expressed as xanthine dehydrogenase (XDH, EC 1.1.1.204), which transfers the electrons to NAD^+ . The final product of XOR




Figure 2. Biologic effects of reactive oxygen species (ROS) and reactive nitrogen species (RNS) and their implication in cancer pathogenesis. Low levels of ROS and RNS may stimulate the inflammatory response, endothelial dysfunction, and cell proliferation, whereas higher levels may induce mutagenesis or have cytotoxic effects. The arrows indicate effects directly (continuous line) or indirectly (dashed line) related to oncogenesis.

activity, uric acid, has an antioxidant action in the extracellular environment, providing protection in the blood circulation against oxidative stress. However, in mammals, XOR may be converted to xanthine oxidase (XO, EC 1.1.3.22) through the oxidation of sulfhydryl residues or limited proteolysis. XO utilizes molecular oxygen as an electron acceptor and produces the reactive oxygen species (ROS) superoxide anion and hydrogen peroxide. XOR can generate ROS also by oxidizing NADH, and can contribute to the formation of reactive nitrogen species (RNS) by converting nitrites into nitric oxide. Both ROS and RNS may have many biologic effects, including endothelial dysfunction, inflammation, induction of proliferation, mutagenesis, and cytotoxicity (reviewed in [4, 5]) (Fig. 2).

Level of Xanthine Oxidoreductase in Cancer Tissue

Activity in animal tumors

The first detection of XOR in neoplastic tissues occurred 80 years ago in the extracts of some rodent tumors [6].

Thereafter, a very low XOR activity was found in rat liver during carcinogenesis induced by p-dimethylaminoazobenzene and a low-protein diet, while a parallel increase in serum enzyme was observed, suggesting a “leakage” of the liver enzyme into the blood stream [7]. After feeding rats for a short period with carcinogenic azo-dyes, a decrease in XOR activity level was somewhat present in precancerous liver and was remarkable in the resulting hepatoma [8]. Diethylnitrosamine by intraperitoneal injection induced

hepatocellular carcinoma in rats. During the oncogenesis process, a significant XOR decrease in hepatic tissue was observed compared with that in liver of untreated rats [9]. Additionally, XOR activity was lower in rapidly growing transplantable hepatocellular carcinoma HC-252 than in resting normal rat liver tissue [10].

The catabolism of purines in rat Novikoff hepatoma was deeply impaired, and no activity of XOR could be detected [11]. Accordingly, in chemically induced transplantable hepatomas in rat, the activities of the anabolic purinic enzymes increased, whereas the activities of catabolic enzymes, including XOR, decreased [12]. This alteration may be functional to cancer cell growth because it allows the recycling of purines through the salvage pathway for nucleic acid biosynthesis [13] (reviewed in [14]).

Xanthine oxidoreductase activity progressively decreases also during carcinogenesis in mouse breast tissue [15]. In particular, the XOR level in breast tissue of a low-tumor strain was twice than in that of a high-tumor strain and was further halved in spontaneous mammary tumors [16].

However, the variation in XOR level is not strictly dependent on cell proliferation, since XOR activity increased markedly in mouse breast during pregnancy and lactation [17, 18], as well as in rat liver during postnatal growth [13, 19]. Also, no changes were reported in XOR activity of regenerating rat liver compared with the liver of sham-operated controls [13, 19]. Moreover, as compared with normal rat liver, a 2- to 10-fold decrease in XOR activity was reported in various hepatoma cell lines, irrespectively of the growth rate or degree of histologic differentiation [13, 19]. At a contrast with the precedent observation, the XOR activity decrement in the rapidly growing Morris hepatoma 3924A was higher than in the slowly growing chemically induced transplantable hepatoma 20, being 4 and 34% of those of normal rat liver, respectively [20].

As compared with the levels in normal intestinal tissue, significantly lower levels of XOR activity have been reported in mouse colon adenocarcinoma [21], as well as in chemically induced tumors of the rat large bowel [22]. XOR activity in the serum of treated rats was considerably higher than that of the control animals [22]. A metabolic imbalance was reported in mouse colon adenocarcinoma favoring purine synthesis over degradation, in a manner similar to that observed in rat hepatomas [21].

Similar biochemical alterations were described in rat kidney tumors [23], as well as in a rapidly growing rat rhabdomyosarcoma, with the level of XOR activity at 54% compared with that in muscle tissue, independently of the rate of cell replication because XOR activity was doubled in a differentiating muscle compared with adult muscle [24].

Contrasting results were obtained with murine skin during carcinogenesis, and in papillomas and squamous cell carcinomas induced by 12-O-tetradecanoylphorbol-13-acetate. The activities of several antioxidant enzymes were reduced, whereas XOR activity was significantly increased, as measured in age-matched, nontreated mice or in skin adjacent to the tumors [25]. These findings are somewhat in agreement with the results obtained with the transplantable Guerin's epithelioma in rats. The level of XOR activity and, in particular its oxidase form, increased during the period of carcinoma active growth together with the formation of ROS. XOR activity declined at the terminal stages of oncogenesis, when the enzyme protein was degraded, suggesting a role for ROS in the mechanism of tumor pathogenesis [26].

In most of the cases, XOR expression and activity decreased in tumor tissues compared with the corresponding normal ones. The experimental reports are generally focused on the liver, breast, colon, and kidney cancers, that is, tumors derived from the tissues with the highest XOR activity. Thus, it is not surprising that a lack of differentiation in malignant neoplasms may correspond to a low level of a highly expressed protein such as XOR. However, in tumors derived from low XOR-expressing tissues, such as skin squamous cell carcinomas, other considerations should be made. For instance, an inflammatory response with an increase in the XOR activity level could be expected.

Activity in human tumors

In two lines of human colon carcinoma xenografts, which were carried in nude mice, purine metabolism appeared to depend on the different growth rates. The rapidly growing carcinoma cells showed a higher level of purine biosynthetic enzymes and a lower level of purine catabolic enzymes compared with the slower growing cell line. Particularly, the rate-limiting XOR activity in the rapidly growing and poorly differentiated colon cancer was decreased to 17% compared with the slower growing, well-differentiated tumor [27]. Additionally, XOR activity decreased, whereas the level of enzymes involved in the purine salvage pathway increased, in cancerous human colon tissues compared with noncancerous ones, conferring selective growth advantages to neoplastic tissues [28].

Xanthine oxidoreductase activity was significantly decreased in primary renal cell carcinomas compared with control kidney cells [23]. Similar alterations of the purine metabolism were observed by comparing cancerous and noncancerous human tissues from patients with bladder cancer [29] or renal cell carcinoma [30], suggesting an increase in salvage pathway activity in tumor cells.

The activity of the dehydrogenase form of XOR was decreased, although its oxidase form was increased, in human liver cancers compared with noncancerous human liver tissues. The free radical defense enzymes were found to be depressed, and the DNA turnover was accelerated in the same samples [31]. Accordingly, with the above findings, XOR activity was significantly lower in neoplastic than in control tissue, as well as in a nontumoral area of the same liver, in patients with hepatocarcinoma [32].

Low levels of XOR and cellular retinol binding protein have been reported in malignant human mammary epithelial cells compared with normal ones. The consequent retinoic acid deficiency could be responsible for the lack of differentiation and may be implicated in mammary carcinogenesis [33].

Dissimilar results were obtained in patients with tumors of different histologic origin. Significantly higher XOR levels were observed in meningioma and astrocytoma from the human brain compared with normal brain tissue [34]. Increased XOR activity was also reported in human laryngeal well-differentiated squamous cell carcinomas compared with the corresponding tumor-free adjacent tissues or normal laryngeal tissues [35].

Increased activities of the enzymes participating in purine metabolism, including XOR activity, have also been described in cancerous human colorectal tissues, whereas the activities of free radical-metabolizing enzymes decreased compared with those of noncancerous adjacent tissues. The increment in XOR activity was interpreted as a reaction to accelerated DNA turnover, while the reduced antioxidant defense could be, at least in part, responsible for the carcinogenic processes [36].

In cancerous human prostate tissue, compared with noncancerous tissues, the activities of DNA turnover enzymes decreased, and the activities of free radical-metabolizing enzymes increased, although the activities of XOR and superoxide dismutase were unchanged. The overall results suggested that a low rate of purine catabolism, as well as an increased free radical stress occur in tumoral prostate tissues [37].

Finally, as sometimes reported in tumor-bearing animals, an elevated level of XOR activity was also observed in the plasma of patients with different types of cancer [38].

Expression in human tumors

Normal human breast has been shown to express XOR protein in the cytoplasm of the epithelium lining terminal ducts. XOR expression was markedly enhanced in the alveolar epithelium of lactating mammary lobules. No detectable XOR was observed either *in situ* or in invasive carcinomas of the breast [39]. In agreement with these results, XOR expression was strong in normal breast

epithelium and was expressed at different levels in 1262 breast cancers, independently from the hormone receptor status of the tumor. A low level of expression was found in large-sized cancers with poor histologic grade of differentiation and a high number of positive axillary lymph nodes. The loss of XOR expression was associated with an unfavorable prognosis. Indeed, the risk of distant recurrence was doubled in patients with no XOR expression compared with those with a moderately decreased or normal expression. Thus, the absence of XOR expression was an independent prognostic factor, associated with unfavorable outcomes. However, dedifferentiation did not invariably lead to loss of XOR [40]. Moreover, the XOR level was found to be higher in breast cancer cell lines with no or low migratory capability than in highly invasive mammary tumor cells, suggesting a potential role for XOR in suppressing breast cancer pathogenesis [41].



Xanthine oxidoreductase expression in 337 human gastric cancers varied from normal to undetectable levels compared with the corresponding normal tissue. Downregulation of XOR expression was associated with the parameters of a more aggressive gastric cancer in terms of cellular anaplasia, mitotic index, invasiveness, tumor size, and lymph node metastasis. Low XOR expression resulted in a marker of unfavorable outcome and poor specific survival in patients with gastric cancer [42].

Similar results were obtained by comparing the level of XOR expression in human colorectal cancer and in normal intestinal tissue of 478 patients. A variable decrease was observed in most of the tumor samples; however, poor XOR expression was associated with a low grade of histologic differentiation and a more advanced cancer stage. Moreover, no XOR expression was detected *in vitro* in the undifferentiated Caco-2 cell line, derived from human colon adenocarcinoma, but XOR expression appeared and increased when these cells spontaneously differentiated during cell culture. Thus, the level of XOR expression may also be a prognostic factor for the clinical outcomes of colorectal carcinomas, suggesting a neoplasia suppressor role for XOR in this tumor, as well as in breast and gastric cancers [43].

Decreased XOR expression in tumor tissue compared with the corresponding normal tissue was associated with a worse prognosis in 474 patients with serous ovarian carcinoma; XOR expression was markedly lower in poorly differentiated tumors. Moreover, the outcome and specific survival were negative, even in cancers with an otherwise more favorable prognostic profile, when XOR was absent [44].

The expression of XOR protein was absent in normal human lung tissue, while it was observed at various levels in 65% of different types of non-small-cell lung cancers. Patients with a high percentage of cells expressing XOR

Tissues normally expressing XOR at a low level



Tissues normally expressing XOR at a high level

Figure 3. Expression and activity of Xanthine oxidoreductase (XOR) in cancer. The level of XOR expression and activity in cancer tissue was down- or upregulated in the malignancies of tissues, which express XOR protein at high (liver, breast, colon, and kidney) or low (skin and lung) levels, respectively.

protein had a longer median survival than patients with a lower expression frequency. Additionally, a low intensity of XOR expression was associated with a worse prognosis for patients who received adjuvant chemotherapy, suggesting a role for XOR-produced ROS in contributing to the tumoricidal effect of chemotherapy [45]. While the lack of XOR expression was confirmed in human lung tissue, the acquisition of such expression was associated with a shortened survival in patients with resected lung adenocarcinoma. Thus, the level of XOR mRNA in tumor tissue was correlated with the worsening of patient outcomes, suggesting it to be an independent predictor of a poor prognosis [46].

A higher variability related to the XOR level was observed in the clinical results compared with the experimental ones. This variability cannot be unexpected for obvious reasons because the laboratory animals have more homogeneous genetic material. However, low XOR expression was associated with a high malignancy grade and a worse

prognosis in patients with breast, liver, gastrointestinal, and kidney cancers, which are derived from tissues expressing a high level of XOR protein. Similar to what was observed in animal tumors, an increased XOR level in low-expressing tissues, as well as the acquired XOR expression in the usually not expressing lung tissue, may have a divergent meaning, possibly indicating an inflammatory reaction, in particular being associated with the aggravation of patient outcome (Fig. 3).


Role of Xanthine Oxidoreductase in Differentiation and Oncogenesis

Development and differentiation

Despite some controversial results, the alteration of XOR expression and activity, both in experimental and clinical reports, suggested a role for XOR in the differentiation process. Indeed, XOR has been implicated in cell differentiation because of its ability to regulate various molecules involved in the transduction of inter- and intracellular signals: (1) pro-inflammatory cyclooxygenase-2 (COX-2) [41, 47], (2) transcription factor NF- κ B (nuclear factor kappa-light-chain-enhancer of activated B cells) [48, 49], (3) hypoxia-inducible factor-1 α (HIF-1 α) [50], and (4) peroxisome proliferator-activated receptor- γ (PPAR- γ) [51] (Fig. 4A).

XOR $+/-$ mice showed defective production and secretion of milk fat droplets that impair the lactation process [52]. Indeed, XOR has been demonstrated to be essential for the regulation of milk lipid formation and secretion in the mouse mammary gland [53]. XOR $-/-$ mice also had a defective induction of COX-2, associated with renal dysplasia, probably because of the uric acid lack [47]. Moreover, these mice had an accumulation of triglycerides with injury of renal tubules and interstitial fibrosis possibly due to increased epithelial–mesenchymal transition [54]. The amount of adipose mass was proportional to XOR expression by comparing XOR $-/-$, wild-type, and obese *ob/ob* mice, accrediting XOR as a regulator of adipogenesis [51]. Additionally, estrogen-dependent XOR expression and the conversion of XDH to XO were associated with the orthokeratinization pathway of differentiation in the vaginal epithelium of ovariectomized mice [55].

In mammary epithelial cell lines with weak XOR expression and high migratory capacity, the overexpression of XOR cDNA reduced COX-2 expression and inhibited cell migration in vitro. Conversely, in cell lines with high XOR expression and weak migratory capacity, XOR inhibition promoted increased levels of COX-2 protein and cell migration in vitro, suggesting a potential role for XOR in suppressing the invasiveness of breast cancer



through the regulation of COX-2 [41]. XOR expression was downregulated by estrogens, which may promote oncogenesis, whereas the differentiation of mammary epithelial cells was associated with XOR activity increase both in vitro and in vivo. Low XOR expression contributed to breast cancer aggressiveness and predicted a more rapid time to tumor relapse. Similarly, the inhibition of XOR increased tumor growth in a mouse xenograft model of human breast cancer [56]. Thus, XOR may be considered a marker of mammary gland development and differentiation (Fig. 4B).

Oncogenesis

The reaction products of XOR activity, uric acid and ROS/RNS have been implicated both in the process of oncogenesis, as well as in its prevention (reviewed in [57]).

The presence of XOR occurs in bladder tissue, and the oxidase form of XOR could catalyze the metabolic activation of a bladder carcinogen, supporting the hypothesis that such activation could occur in vivo [58]. Accordingly, the colon tumorigenesis, induced in rats by azo-dye feeding, was prevented by the oral administration of a XOR inhibitor [59].

Alcohol intake is associated with an increased risk of breast cancer, and a pathogenetic role has been suggested for XOR because acetaldehyde and NADH, both produced by the metabolism of alcohol, can be substrates for XOR, resulting in ROS formation. Moreover, the age-induced

Figure 4. Molecular and cellular actions of Xanthine oxidoreductase (XOR). (A) Signal transduction molecules upregulated by XOR. COX-2 is an inducible enzyme producing pro-inflammatory metabolites of arachidonic acid, which are involved in oncogenesis and progression to malignancy by promoting cell proliferation and migration. COX-2 expression requires the phosphorylation of NF-κB and could be inhibited by XOR [41], but stimulated by uric acid [47]. NF-κB is a transcription factor that may be activated by XOR-produced reactive oxygen species (ROS) [48, 49]. NF-κB activity is a hallmark of chronic inflammation and cancer, because its signaling pathway maintains high expression of a set of antiapoptotic genes. HIF-1α is a transcription factor implicated in cancer development and progression by promoting the expression of angiogenic target genes, including vascular endothelial growth factor. In normoxic conditions, activated NF-κB contributes to the expression and activity of HIF-1α together with ROS, including XOR-derived ROS [50]. Peroxisome proliferator-activated receptor gamma (PPAR)-γ is a ligand-activated intracellular transcription factor, included in the nuclear hormone receptor superfamily that has an antitumorigenic and antiproliferative action by favoring cell differentiation and inhibiting angiogenesis. XOR stimulated the activation of PPAR-γ through the ROS produced by its NADH-oxidizing activity [51]. (B) Cell differentiation aspects affected by XOR. Milk fat droplet production and secretion requires the formation of a complex between XOR, adipophilin and butyrophilin [53]. Thus, in lactating mammary epithelium of mice, XOR expression is essential to maintain lactation [52], possibly because XOR regulates the binding of fat droplets to the apical cell membrane and their secretion [18]. Pluristratified epithelium orthokeratinization was associated to XOR expression and conversion of xanthine dehydrogenase (XDH) to xanthine oxidase (XO) in ovariectomized mice after hormonal induction of vaginal epithelium hyperplasia, suggesting that XOR activity is required for the orthokeratinization pathway of differentiation [55]. Adipogenesis is controlled by a cascade of factors that includes XOR as a regulator of the adipogenic process. Indeed, in vivo the level of XOR expression correlates with fat accumulation in mice and XOR activity was linked to the adipogenic transcription factor PPAR-γ [51]. Epithelial to mesenchymal transition was observed in primary renal epithelial cells from XOR-/- mice [54]. Not only the cell morphology changed from a cuboidal to a myofibroblastic shape, but also immunohistochemical analysis showed a positive staining for markers of mesenchymal cell type, suggesting that XOR expression is needed for epithelial differentiation in mouse kidney. Migratory capacity in vitro was inversely correlated with XOR expression in epithelial cell lines derived from breast cancer [41].

diminished antioxidant defenses and iron accumulation in breast tissue may increase ROS production with the risk of DNA damage and carcinogenic mutations (reviewed in [60]). Such in situ bioactivation was shown in breast tissue from postlactation rats, demonstrating the ability of XOR to catalyze ethanol to acetaldehyde and free radicals, which may exert a transforming action [61]. The role of XOR-derived ROS in oncogenesis has been suggested by in vitro study, showing that *hXOR* is targeted by tumor suppressors which repress *hXOR* expression by binding to its promoter [62]. The XOR mRNA levels are normally low in most of the tissues because of the repressing regulation at the transcriptional level [63].

A low activity of various oxidative enzymes, including XOR, was correlated with cell proliferation, and the proposed hypothesis was that a low level of free oxygen radicals may promote cancer cell growth [10]. However, ROS-mediated NF κ B activation was suggested to increase the expression of genes involved in angiogenesis and metastasis in human cancer [64]. XOR-produced ROS upregulated HIF-1 α expression [49]. HIF-1 α together with activated NF κ B, contributes to cancer-associated inflammatory signaling, as well as to tumor progression by activating angiogenesis, migration, and invasion (reviewed in [65]).

In intestinal epithelial cells, phlogistic NF- κ B signaling can induce dedifferentiation into tumor-initiating cells. The chemical and biological mechanisms involved in the etiology of inflammation-induced colon cancer were analyzed in correlation with the availability of ROS and RNS. The hypothesis was formulated that these oxidants, produced during inflammation by various sources, including phagocyte XOR, induce oxidative damage to DNA, generating base lesions, strand breaks, genomic aberrations, and chromosomal rearrangements. As a consequence, XOR products can contribute to genetic and epigenetic changes leading to mutation, tumor initiation, and promotion that are responsible for neoplastic transformation through activation of oncogenes and/or inactivation of tumor suppressors (reviewed in [66]).

Controversial role of uric acid

High serum uric acid levels were associated with a short survival time in 118 terminally ill patients with cancer in various organs, including lung and abdominal viscera as primary sites and liver, lung, bone, and brain as metastatic sites [67]. Additionally, the serum uric acid level and XOR activity were markedly increased in patients during hepatocarcinoma progression from the lower to higher stages [68]. Accordingly, the inhibition of XOR activity improved the outcome in a rat model of cancer cachexia, suggesting a role for XOR-produced ROS in inflammatory signaling and tissue wasting [69]. Thus, both an increment in cell damage and a deterioration of renal function could be responsible for hyperuricemia in end-stage cancer patients.

Uric acid lowered E-cadherin levels in cultured NRK renal tubular cells either by decreasing its synthesis and increasing its degradation. Experimental hyperuricemia in rats was associated with the downregulation of E-cadherin expression and increase in alpha-smooth muscle actin expression, indicating an epithelial–mesenchymal transition. These phenotypic alterations, which occur during tumor progression toward malignity, were inhibited by allopurinol [70].

The plasma level of uric acid was significantly higher in patients with squamous cell carcinoma of the head and neck region than in control subjects. Uricemia was also observed to increase with staging, thus behaving as a diagnostic and prognostic marker of head and neck cancers [71]. Hyperuricemia may contribute to cancer pathogenesis by promoting an increased intracellular concentration of uric acid, with consequent pro-inflammatory effects due to ROS and RNS generation and COX-2 activation. After the neoplastic transformation, the scavenger action of extracellular uric acid may protect cancer cells from oxidative stress, thus contributing to their survival (reviewed in [72]). In addition, the uric acid can reduce the tumor surveillance by the immune system as it is able to stimulate the chemotaxis of mesenchymal stromal human cells, which are associated with tumor progression and metastasis and can promote the differentiation of immunosuppressive IDO+/IL10+ lymphocytes [73]. Accordingly, an elevated uric acid level in plasma at the time of diagnosis was a negative prognostic marker in a cohort of 466 pancreatic cancer patients [74]. Also, in two cohort studies including 83,683 men and 28,613 women with a median follow-up of 13.6 and 15.2 years, respectively, high serum uric acid concentrations were associated with increased risk of cancer mortality [75, 76]. Moreover, the above wide cohort of male patients was further analyzed, investigating the association of serum uric acid and cancer incidence, by using a nonlinear statistical test (penalized splines in extended Cox-type additive hazard regression). The results of the study evidenced the “J-shaped” nature of the relationship between serum uric acid and cancer incidence and indicated a positive dose–response association between these parameters for serum uric acid above the normal level [77].

In contrast, the antioxidant activity of uric acid could exert a contrasting action against free-radical oncogenesis. The incidence of lung cancer (1015 cases) in cigarette smokers was inversely related to the serum uric acid level [78]. The hypothesis has been formulated that the loss of uricase during primate evolution may have had a crucial role in lengthening life-span because of the scavenger activity of uric acid (reviewed in [79]) with the consequent reduction in age-specific cancer rates [80]. A link has been demonstrated between the tumor suppression by p53 and antioxidant function of uric acid through its transporter SLC2A9, whose expression is induced by oxidative stress and is dependent on p53. An increased ROS level was reported by inhibiting SLC2A9 with small-interfering RNA or because of p53 loss. Thus, the poorer prognosis associated with decreased SLC2A9 expression that was described in several cancer types suggests that the p53–SLC2A9 system could prevent the ROS-associated damage contributing to cancer development [81].

Elevated levels of uric acid have been associated with better survival in various malignancies. In colorectal cancer patients, a better prognosis was associated with an elevated level of serum uric acid, suggesting this as an efficient predictor of survival [82]. The prognostic relevance of preoperative uric acid plasma levels to cancer-specific survival was evaluated in 357 soft-tissue sarcoma patients who underwent tumor resection. Conclusions similar to those above reported were suggested by the positive clinical outcome, which was associated with elevated uric acid serum levels [83]. The protection against cancer mortality given by elevated serum uric acid levels was demonstrated by a large general population-based cohort study with 4350 patients who died because of cancer, in particular, cancer localized to the lung, colorectal intestine, and prostate [84].

Conclusion

In most tissues, *hXOR* is normally downregulated at the transcriptional level, except in the breast, liver, gastrointestinal tract, and kidney. Differentiation is associated with the expression of a high level of XOR protein in these organs, whereas the lack of differentiation, which occurs in malignant neoplasms, corresponds to a low concentration of XOR. In addition, in the case of the mammary gland, XOR plays a structural role and may be considered a marker of development and differentiation. By activating PPAR- γ , XOR promotes cell differentiation and produces an antitumorogenic and antiproliferative action. Moreover, the low XOR activity, together with an increase in purine biosynthesis, provides a selective advantage to the growth of tumor tissue. Indeed, XOR expression was negatively associated with a high malignancy grade and a worse prognosis in cancer patients.

The elevated XOR activity level observed in the plasma of patients with different types of cancer may be related to the inflammatory response elicited by the tissue damage induced by tumor growth. A similar reaction is possibly responsible for an elevation of XOR expression in usually low-expressing tissues; if this interpretation is correct, the level of XOR expression should be positively associated with a worse outcome in these cancer patients.

The role of XOR activity and its products in oncogenesis has been suggested. XOR in its oxidase form could directly catalyze the metabolic activation of carcinogenic substances. In most cases, XOR has an indirect tumorigenic action by producing ROS and RNS, which may activate proinflammatory, antiapoptotic, and angiogenic target genes, by NF- κ B and HIF-1 α upregulation. Uric acid promotes inflammation, cell proliferation, and migration by inducing COX-2 expression. However, the role of uric acid in cancer is still debated. The antioxidant function of circulating uric acid justifies the overall protective role against

neoplasia, although the elevation of its plasma level is typical of the end stages of oncologic illness that is associated with tumor lysis syndrome.

Current knowledge does not allow to suggest the use of available XOR inhibitors to prevent both ROS/RNS formation and oncogenesis. However, XOR inhibitors have a wide pharmacological use in hyperuricemic patients with metabolic syndrome, renal diseases or tumor lysis syndrome. Thus, a large amount of data will be available in the future for further studies to understand the correlation between XOR activity, reflected by the serum uric acid level, and cancer incidence and prognosis. The results of such meta-analysis will possibly discriminate the cases in which the control of the uricemia will be helpful to improve the outcomes of tumor illness. In addition, a conditional knockout strategy in mice would be useful to test the differential role of XOR and uric acid in pre-clinical model of tumorigenesis and cancer progression.

Acknowledgments

This work was supported by the Pallotti Legacies for Cancer Research. No relationship is present regarding any organization or entity having a direct financial or personal interest in the subject matter or materials discussed in the article.

Conflict of Interest

None declared.

References

1. Pritsos, C. A., and D. L. Gustafson. 1994. Xanthine dehydrogenase and its role in cancer chemotherapy. *Oncol. Res.* 6:477–481.
2. Hille, R., and T. Nishino. 1995. Flavoprotein structure and mechanism. 4. Xanthine oxidase and xanthine dehydrogenase. *FASEB J.* 9:995–1003.
3. Chung, H. Y., B. S. Baek, S. H. Song, M. S. Kim, J. I. Huh, K. H. Shim, et al. 1997. Xanthine dehydrogenase/xanthine oxidase and oxidative stress. *Age (Omaha)* 20:127–140.
4. Battelli, M. G., A. Bolognesi, and L. Polito. 2014a. Pathophysiology of circulating xanthine oxidoreductase: new emerging roles for a multi-tasking enzyme. *Biochim. Biophys. Acta* 1842:1502–1517.
5. Battelli, M. G., L. Polito, and A. Bolognesi. 2014b. Xanthine oxidoreductase in atherosclerosis pathogenesis: not only oxidative stress. *Atherosclerosis* 237:562–567.
6. Boyland, E., and M. E. Boyland. 1935. Studies in tissue metabolism: lactic dehydrogenase, xanthine oxidase and nucleosidase in tumour and muscle extracts. *Biochem. J.* 29:1097–1101.

7. Westerfeld, W. W., D. A. Richert, and M. F. Hilfinger. 1950. Studies on xanthine oxidase during carcinogenesis by p-dimethylaminoazobenzene. *Cancer Res.* 10:486–494.
8. Reid, E., and I. Lewin. 1957. Adenosine deaminase, nucleoside phosphorylase and xanthine oxidase in liver tumours. *Br. J. Cancer* 11:494–498.
9. Metwally, N. S., T. E. Kholeif, K. Z. Ghanem, A. R. Farrag, N. M. Ammar, and A. H. Abdel-Hamid. 2011. The protective effects of fish oil and artichoke on hepatocellular carcinoma in rats. *Eur. Rev. Med. Pharmacol. Sci.* 15:1429–1444.
10. Sun, A. S., and A. I. Cederbaum. 1980. Oxidoreductase activities in normal rat liver, tumor-bearing rat liver, and hepatoma HC-252. *Cancer Res.* 40:4677–4681.
11. de Lamirande, G., C. Allard, and A. Cantero. 1958. Purine-metabolizing enzymes in normal rat liver and Novikoff hepatoma. *Cancer Res.* 18:952–958.
12. Weber, G. 1983a. Enzymes of purine metabolism in cancer. *Clin. Biochem.* 16:57–63.
13. Prajda, N., H. P. Morris, and G. Weber. 1976. Imbalance of purine metabolism in hepatomas of different growth rates as expressed in behavior of xanthine oxidase (EC 1.2.3.2). *Cancer Res.* 36:4639–4646.
14. Weber, G. 1983b. Biochemical strategy of cancer cells and the design of chemotherapy: G. H. A. Clowes Memorial Lecture. *Cancer Res.* 43:3466–3492.
15. Sheth, N. A., S. V. Bhide, and K. J. Ranadive. 1968. Xanthine oxidase activity in progressive spontaneous mammary carcinogenesis. *Br. J. Cancer* 22:833–838.
16. Lewin, I., R. Lewin, and R. C. Bray. 1957. Xanthine oxidase activity during mammary carcinogenesis in mice. *Nature* 180:763–764.
17. Lewin, I. 1957. Discussion on mammary tumors: xanthine oxidase activity in normal & abnormal growth. *Proc. R. Soc. Med.* 50:563–570.
18. McManaman, J. L., C. A. Palmer, R. M. Wright, and M. C. Neville. 2002. Functional regulation of xanthine oxidoreductase expression and localization in the mouse mammary gland: evidence of a role in lipid secretion. *J. Physiol.* 545:567–579.
19. Prajda, N., and G. Weber. 1975. Malignant transformation-linked imbalance: decreased xanthine oxidase activity in hepatomas. *FEBS Lett.* 59:245–249.
20. Ikegami, T., Y. Natsumeda, and G. Weber. 1986. Decreased concentration of xanthine dehydrogenase (EC 1.1.1.204) in rat hepatomas. *Cancer Res.* 46:3838–3841.
21. Weber, G., H. Kizaki, D. Tzeng, T. Shiotani, and E. Olah. 1978. Colon tumor: enzymology of the neoplastic program. *Life Sci.* 23:729–736.
22. Mitidieri, E., A. S. Souza, and O. R. Affonso. 1988. Xanthine oxidase activity in intestinal tumors induced by 1,2-dimethylhydrazine dihydrochloride. *Arch. Geschwulstforsch* 58:79–80.
23. Prajda, N., J. P. Donohue, and G. Weber. 1981. Increased amidophosphoribosyltransferase and decreased xanthine oxidase activity in human and rat renal cell carcinoma. *Life Sci.* 29:853–860.
24. Weber, G., M. E. Burt, R. C. Jackson, N. Prajda, M. S. Lui, and E. Takeda. 1983. Purine and pyrimidine enzymic programs and nucleotide pattern in sarcoma. *Cancer Res.* 43:1019–1023.
25. Reiners, J. J. Jr, G. Thai, T. Rupp, and A. R. Cantu. 1991a. Assessment of the antioxidant/prooxidant status of murine skin following topical treatment with 12-O-tetradecanoylphorbol-13-acetate and throughout the ontogeny of skin cancer. Part I: quantitation of superoxide dismutase, catalase, glutathione peroxidase and xanthine oxidase. *Carcinogenesis* 12:2337–2343.
26. Shmarakov, I. A., and M. M. Marchenko. 2009. Xanthine oxidase activity in transplantable Guerin's carcinoma in rats. *Vopr. Onkol.* 55:345–350.
27. Weber, G., J. C. Hager, M. S. Lui, N. Prajda, D. Y. Tzeng, R. C. Jackson, et al. 1981. Biochemical programs of slowly and rapidly growing human colon carcinoma xenografts. *Cancer Res.* 41:854–859.
28. Durak, I., R. Cetin, E. Devrim, and I. B. Ergüder. 2005. Effects of black grape extract on activities of DNA turn-over enzymes in cancerous and non cancerous human colon tissues. *Life Sci.* 76:2995–3000.
29. Durak, I., H. Perk, M. Kavutçu, O. Canbolat, O. Akyol, and Y. Bedük. 1994. Adenosine deaminase, 5' nucleotidase, xanthine oxidase, superoxide dismutase, and catalase activities in cancerous and noncancerous human bladder tissues. *Free Radic. Biol. Med.* 16:825–831.
30. Durak, I., Y. Bedük, M. Kavutcu, O. Süzer, O. Yaman, H. S. Oztürk, et al. 1997. Activity of the enzymes participating in purine metabolism of cancerous and noncancerous human kidney tissues. *Cancer Invest.* 15:212–216.
31. Durak, I., M. Karaayvaz, H. S. Ozturk, M. Kacmaz, and H. Akgul. 1998. Activities of DNA turn-over and free radical-metabolizing enzymes and levels of peroxidation indexes in human hepatic cancer tissues. *Cancer Res. Ther. Control* 5:195–201.
32. Stirpe, F., M. Ravaioli, M. G. Battelli, S. Musiani, and G. L. Grazi. 2002. Xanthine oxidoreductase activity in human liver disease. *Am. J. Gastroenterol.* 97:2079–2085.
33. Taibi, G., G. Carruba, L. Cocciadiferro, O. M. Granata, and C. M. Nicotra. 2009. Low levels of both xanthine dehydrogenase and cellular retinol binding protein are responsible for retinoic acid deficiency in malignant human mammary epithelial cells. *Ann. N. Y. Acad. Sci.* 1155:268–272.
34. Kökoğlu, E., A. Belce, E. Ozyurt, and Z. Tepeler. 1990. Xanthine oxidase levels in human brain tumors. *Cancer Lett.* 50:179–181.

35. Durak, I., A. C. Işık, O. Canbolat, O. Akyol, and M. Kavutçu. 1993. Adenosine deaminase, 5' nucleotidase, xanthine oxidase, superoxide dismutase, and catalase activities in cancerous and noncancerous human laryngeal tissues. *Free Radic. Biol. Med.* 15:681–684.
36. Oztürk, H. S., M. Karaayvaz, M. Kaçmaz, M. Kavutcu, H. Akgül, and I. Durak. 1998. Activities of the enzymes participating in purine and free-radical metabolism in cancerous human colorectal tissues. *Cancer Biochem. Biophys.* 16:157–168.
37. Biri, H., H. S. Oztürk, M. Kaçmaz, K. Karaca, H. Tokuçoğlu, and I. Durak. 1999. Activities of DNA turnover and free radical metabolizing enzymes in cancerous human prostate tissue. *Cancer Invest.* 17:314–319.
38. Samra, Z. Q., S. Pervaiz, S. Shaheen, N. Dar, and M. A. Athar. 2011. Determination of oxygen derived free radicals producer (xanthine oxidase) and scavenger (paraoxonase1) enzymes and lipid parameters in different cancer patients. *Clin. Lab.* 57:741–747.
39. Cook, W. S., R. Chu, M. Saksela, K. O. Raivio, and A. V. Yeldandi. 1997. Differential immunohistochemical localization of xanthine oxidase in normal and neoplastic human breast epithelium. *Int. J. Oncol.* 11:1013–1017.
40. Linder, N., J. Lundin, J. Isola, M. Lundin, K. O. Raivio, and H. Joensuu. 2005. Down-regulated xanthine oxidoreductase is a feature of aggressive breast cancer. *Clin. Cancer Res.* 11:4372–4381.
41. Fini, M. A., D. Orchard-Webb, B. Kosmider, J. D. Amon, R. Kelland, G. Shibao, et al. 2008. Migratory activity of human breast cancer cells is modulated by differential expression of xanthine oxidoreductase. *J. Cell. Biochem.* 105:1008–1026.
42. Linder, N., C. Haglund, M. Lundin, S. Nordling, A. Ristimäki, A. Kokkola, et al. 2006. Decreased xanthine oxidoreductase is a predictor of poor prognosis in early-stage gastric cancer. *J. Clin. Pathol.* 59:965–971.
43. Linder, N., E. Martelin, M. Lundin, J. Louhimo, S. Nordling, C. Haglund, et al. 2009. Xanthine oxidoreductase - clinical significance in colorectal cancer and in vitro expression of the protein in human colon cancer cells. *Eur. J. Cancer* 45:648–655.
44. Linder, N., R. Bützow, H. Lassus, M. Lundin, and J. Lundin. 2012. Decreased xanthine oxidoreductase (XOR) is associated with a worse prognosis in patients with serous ovarian carcinoma. *Gynecol. Oncol.* 124:311–318.
45. Kim, A. W., M. Batus, R. Myint, M. J. Fidler, S. Basu, P. Bonomi, et al. 2011. Prognostic value of xanthine oxidoreductase expression in patients with non-small cell lung cancer. *Lung Cancer* 71:186–190.
46. Konno, H., Y. Minamiya, H. Saito, K. Imai, Y. Kawaharada, S. Motoyama, et al. 2012. Acquired xanthine dehydrogenase expression shortens survival in patients with resected adenocarcinoma of lung. *Tumour Biol.* 33:1727–1732.
47. Ohtsubo, T., I. I. Rovira, M. F. Starost, C. Liu, and T. Finkel. 2004. Xanthine oxidoreductase is an endogenous regulator of cyclooxygenase-2. *Circ. Res.* 95:1118–1124.
48. Matsui, N., I. Satsuki, Y. Morita, K. Inaizumi, K. Kasajima, R. Kanoh, et al. 2000. Xanthine oxidase-derived reactive oxygen species activate nuclear factor kappa B during hepatic ischemia in rats. *Jpn. J. Pharmacol.* 84:363–366.
49. Romagnoli, M., M. C. Gomez-Cabrera, M. G. Perrelli, F. Biasi, F. V. Pallardó, J. Sastre, et al. 2010. Xanthine oxidase-induced oxidative stress causes activation of NF-kappaB and inflammation in the liver of type I diabetic rats. *Free Radic. Biol. Med.* 49:171–177.
50. Griguer, C. E., C. R. Oliva, E. E. Kelley, G. I. Giles, J. R. Jr Lancaster, and G. Y. Gillespie. 2006. Xanthine oxidase-dependent regulation of hypoxia-inducible factor in cancer cells. *Cancer Res.* 66:2257–2263.
51. Cheung, K. J., I. Tzamelis, P. Pissios, I. Rovira, O. Gavrilova, T. Ohtsubo, et al. 2007. Xanthine oxidoreductase is a regulator of adipogenesis and PPARgamma activity. *Cell Metab.* 5:115–128.
52. Vorbach, C., A. Scriven, and M. R. Capecchi. 2002. The housekeeping gene xanthine oxidoreductase is necessary for milk fat droplet enveloping and secretion: gene sharing in the lactating mammary gland. *Genes Dev.* 16:3223–3235.
53. McManaman, J. L., C. A. Palmer, S. Anderson, K. Schwertfeger, and M. C. Neville. 2004. Regulation of milk lipid formation and secretion in the mouse mammary gland. *Adv. Exp. Med. Biol.* 554:263–279.
54. Ohtsubo, T., K. Matsumura, K. Sakagami, K. Fujii, K. Tsuruya, H. Noguchi, et al. 2009. Xanthine oxidoreductase depletion induces renal interstitial fibrosis through aberrant lipid and purine accumulation in renal tubules. *Hypertension* 54:868–876.
55. Reiners, J. J. Jr, T. Rupp, and C. J. Conti. 1991b. Modulation of xanthine dehydrogenase and oxidase activities during the hormonal induction of vaginal epithelial differentiation in ovariectomized mice. *Differ.* 47:69–75.
56. Fini, M. A., J. Monks, S. M. Farabaugh, and R. M. Wright. 2011. Contribution of xanthine oxidoreductase to mammary epithelial and breast cancer cell differentiation in part modulates inhibitor of differentiation-1. *Mol. Cancer Res.* 9:1242–1254.
57. Battelli, M. G., L. Polito, M. Bortolotti, and A. Bolognesi. 2015. Xanthine oxidoreductase-derived reactive species: physiological and pathological effects. *Oxid. Med. Cell. Longev.* <http://www.hindawi.com/journals/omcl/aip/394245/>.

58. Swaminathan, S., and J. F. Hatcher. 1986. Xanthine oxidase-mediated mutagenicity of the bladder carcinogen 4-nitrobiphenyl. *Mutat. Res.* 172:37–45.
59. Tanaka, T., K. Kawabata, M. Kakumoto, H. Makita, K. Matsunaga, H. Mori, et al. 1997. Chemoprevention of azoxymethane-induced rat colon carcinogenesis by a xanthine oxidase inhibitor, 1'-acetoxychavicol acetate. *Jpn. J. Cancer Res.* 88:821–830.
60. Wright, R. M., J. L. McManaman, and J. E. Repine. 1999. Alcohol-induced breast cancer: a proposed mechanism. *Free Radic. Biol. Med.* 26:348–354.
61. Castro, G. D., A. M. Delgado de Layño, M. H. Costantini, and J. A. Castro. 2001. Cytosolic xanthine oxidoreductase mediated bioactivation of ethanol to acetaldehyde and free radicals in rat breast tissue. Its potential role in alcohol-promoted mammary cancer. *Toxicology* 160:11–18.
62. Lin, J., P. Xu, P. LaVallee, and J. R. Hoidal. 2008. Identification of proteins binding to E-Box/Ku86 sites and function of the tumor suppressor SAFB1 in transcriptional regulation of the human xanthine oxidoreductase gene. *J. Biol. Chem.* 283:29681–29689.
63. Xu, P., P. LaVallee, and J. R. Hoidal. 2000. Repressed expression of the human xanthine oxidoreductase gene. E-box and TATA-like elements restrict ground state transcriptional activity. *J. Biol. Chem.* 275:5918–5926.
64. Tabata, S., R. Ikeda, M. Yamamoto, S. Shimaoka, N. Mukaida, Y. Takeda, et al. 2014. Thymidine phosphorylase activates NFκB and stimulates the expression of angiogenic and metastatic factors in human cancer cells. *Oncotarget* 5:10473–10485.
65. Balamurugan, K. 2015. HIF-1 at the crossroads of hypoxia, inflammation, and cancer. *Int. J. Cancer.* doi:10.1002/ijc.29519. [Epub ahead of print]
66. Mangerich, A., P. C. Dedon, J. G. Fox, S. R. Tannenbaum, and G. N. Wogan. 2013. Chemistry meets biology in colitis-associated carcinogenesis. *Free Radic. Res.* 47:958–986.
67. Shin, H. S., H. R. Lee, D. C. Lee, J. Y. Shim, K. H. Cho, and S. Y. Suh. 2006. Uric acid as a prognostic factor for survival time: a prospective cohort study of terminally ill cancer patients. *J. Pain Symptom Manage.* 31:493–501.
68. Lin, C. C., and M. C. Yin. 2007. B vitamins deficiency and decreased anti-oxidative state in patients with liver cancer. *Eur. J. Nutr.* 46:293–299.
69. Springer, J., A. Tschirner, K. Hartman, S. Palus, E. K. Wirth, S. B. Ruis, et al. 2012. Inhibition of xanthine oxidase reduces wasting and improves outcome in a rat model of cancer cachexia. *Int. J. Cancer* 131:2187–2196.
70. Ryu, E. S., M. J. Kim, H. S. Shin, Y. H. Jang, H. S. Choi, I. Jo, et al. 2013. Uric acid-induced phenotypic transition of renal tubular cells as a novel mechanism of chronic kidney disease. *Am. J. Physiol. Renal. Physiol.* 304:F471–F480.
71. Dhankhar, R., K. Dahiya, T. K. Sharma, V. S. Ghalaut, R. Atri, and V. Kaushal. 2011. Diagnostic significance of adenosine deaminase, uric acid and C-reactive protein levels in patients of head and neck carcinoma. *Clin. Lab.* 57:795–798.
72. Fini, M. A., A. Elias, R. J. Johnson, and R. M. Wright. 2012. Contribution of uric acid to cancer risk, recurrence, and mortality. *Clin. Transl. Med.* 1:1–16.
73. Eisenbacher, J. L., H. Schrezenmeier, B. Jahrsdörfer, C. Kaltenmeier, M. T. Rojewski, T. Yildiz, et al. 2014. S100A4 and uric acid promote mesenchymal stromal cell induction of IL-10 + /IDO+ lymphocytes. *J. Immunol.* 192:6102–6110.
74. Stotz, M., J. Szkandera, J. Seidel, T. Stojakovic, H. Samonigg, D. Reitz, et al. 2014. Evaluation of uric acid as a prognostic blood-based marker in a large cohort of pancreatic cancer patients. *PLoS ONE* 9:e104730.
75. Strasak, A. M., K. Rapp, W. Hilbe, W. Oberaigner, E. Ruttmann, H. Concin, et al. 2007a. Serum uric acid and risk of: cancer mortality in a large prospective male cohort. *Cancer Causes Control* 18:1021–1029.
76. Strasak, A. M., K. Rapp, W. Hilbe, W. Oberaigner, E. Ruttmann, H. Concin, et al. 2007b. The role of serum uric acid as an antioxidant protecting against cancer: prospective study in more than 28000 older Austrian women. *Ann. Oncol.* 18:1893–1897.
77. Strasak, A. M., S. Lang, T. Kneib, L. J. Brant, J. Klenk, W. Hilbe, et al. 2009. Use of penalized splines in extended Cox-type additive hazard regression to flexibly estimate the effect of time-varying serum uric acid on risk of cancer incidence: a prospective, population-based study in 78,850 men. *Ann. Epidemiol.* 19:15–24.
78. Horsfall, L. J., I. Nazareth, and I. Petersen. 2014. Serum uric acid and the risk of respiratory disease: a population-based cohort study. *Thorax* 69:1021–1026.
79. Cutler, R. G. 1991. Antioxidants and aging. *Am. J. Clin. Nutr.* 53:3738–3793.
80. Ames, B. N., R. Cathcart, E. Schwiers, and P. Hochstein. 1981. Uric acid provides an antioxidant defense in humans against oxidant- and radical-caused aging and cancer: a hypothesis. *Proc. Natl Acad. Sci. USA* 78:6858–6862.
81. Itahana, Y., R. Han, S. Barbier, Z. Lei, S. Rozen, and K. Itahana. 2015. The uric acid transporter SLC2A9 is a direct target gene of the tumor suppressor p53 contributing to antioxidant defense. *Oncogene* 34:1799–1810.
82. Dziaman, T., Z. Banaszekiewicz, K. Roszkowski, D. Gackowski, E. Wisniewska, R. Rozalski, et al. 2014.

- 8-Oxo-7,8-dihydroguanine and uric acid as efficient predictors of survival in colon cancer patients. *Int. J. Cancer* 134:376–383.
83. Szkandera, J., A. Gerger, B. Liegl-Atzwanger, M. Stotz, H. Samonigg, F. Ploner, et al. 2015. Uric acid levels in blood are associated with clinical outcome in soft-tissue sarcoma patients. *Clin. Chem. Lab. Med.* 53:493–497.
84. Taghizadeh, N., J. M. Vonk, and H. M. Boezen. 2014. Serum uric acid levels and cancer mortality risk among males in a large general population-based cohort study. *Cancer Causes Control* 25:1075–1080.